

Thermodynamic and Physicochemical Investigation of Rubidium Soaps in Alkanols

S.P. S. Saroha¹, K. Mehrotra², D.Ganjewala¹

¹ Amity Institute of Biotechnology, Amity University, 125, Noida, Uttar Pradesh, India

² Department of Chemistry, University of Jodhpur, Jodhpur, Rajasthan, India

Abstract

Imperative to evolve thermodynamic characteristic of Rubidium carboxylate (caprylate, caprate and laurate) apparent molar volume insight from density studies have been measured in alkanols. Limiting apparent molar volume and experimental slopes S_v too are determined from Masson equation $\phi_v = \phi_{v0} + s_v$. ϕ_{v0} has been interpreted in terms of solute-solvent interaction. Ion-solvent interaction and solvent-solvent interaction are quite stronger in alkanols an insight in the structure making or breaking capacity of individual ions.

Key Words: Apparent molar volume, Caprylate, Capacity, Density, Interaction, Structure, Thermodynamics

INTRODUCTION

Amphiphilic carboxylates of alkali, alkaline earth and transitional metals are of special molecular structure effective for readily association of surfactant molecules naturally find enormous applications in industry and medicinal area. In this venture metal alkanoates of dicarboxylic acids were synthesized and investigated to reveal their ability for the stability of PVC [1-4]. The molecular inter-action of metal alkanoates [5-9] in non- aqueous solvent has crucial role for their application as stabilizer, fungicide, pesticide, foaming, wetting, lubricating, grease, catalyst and emulsifier. The metal carboxylates are characterized [10-12] by physico-chemical studies which give insight to their structure valuable for biological processes. In this paper, we have tried to get information about physical characters specially interaction between anionic head group and metal ion, solute- solvent inter-action, solvent- solvent inter action, apparent molar volume of rubidium soaps in alkanols may be relevant for material sciences.

METHODOLOGY

The materials used rubidium carbonate for synthesis were of BDH/AR grade, the fatty acids obtained from commercial sample were distilled under reduced pressure and further purified by recrystallization was confirmed by the determination of their melting points. Rubidium Carbonate (AR) corresponding to 0.01M was weighted and dissolved in water and the solution was heated to 50-60°C. Requisite amount of fatty acid necessary for complete conversion of rubidium carbonate to soap (0.01M) was melted and added to warm solution of rubidium carbonate with constant stirring. The soap so obtained was digested on water bath for about 2 hours till evolution of carbon dioxide ceases. The excess of acid was removed by washing with benzene and the soaps were purified by recrystallization. After initial drying in an air oven at 100-105°C, the final drying was carried out under reduced pressure and characterized by IR spectra, elemental analysis as in previous article [13].

RESULT AND DISCUSSION

Critical micelle concentration CMC of rubidium soaps in alkanols is obtained from the calculated values of apparent molar volume using following equation:

$$\phi_v = M/d_0 - 1000(d-d_0)/d_0 c \dots\dots\dots (1)$$

Where ϕ_v is apparent molar volume (solute-solvent interaction), M is the molecular mass of soap, d and d_0 are the density of solvent and solution respectively and C is the molar concentration. The interaction of solute and solvent mostly in amphiphilic soaps brings a

change in the volume of solution; this along with the volume of solute is known as apparent molar volume. The apparent molar volume is a linear function of the square root of molar concentration Fig.1 (A, B, C). Plot of apparent molar volume and the square root of molar concentration are linear, show a break at definite soap concentration critical micelle concentration CMC Table 1A.

Table 1A: Critical micelle concentration from graph of ϕ_v vs \sqrt{C} g mol L⁻¹

Soap	Critical micelle concentration		
	Methanol	Butanol	Pentanol
Caprylate	0.044	0.045	0.045
Caprate	0.045	0.046	0.045
Laurate	0.044	0.045	0.044




Figure 1C: Apparent molar volume ϕ_v cm³mol⁻¹ vs square root of concentration g mol L⁻¹




Figure 1 A: Apparent molar volume ϕ_v cm³mol⁻¹ vs square root of concentration g mol L⁻¹




Figure 1 B: Apparent molar volume ϕ_v cm³mol⁻¹ vs square root of concentration g mol L⁻¹

The value of CMC is independent of hydrocarbon chain in soaps. The existence of polar head group of soap molecule causes change in the energy of solution and micellization takes place. It can be argued that a change in overall structure cannot be over ruled.

Massion equation $\phi_v = \phi_{v0} + S_v \sqrt{C}$ is applicable to curve ϕ_v vs \sqrt{C} .

The value of ϕ_{v0} is obtained from the intercept of the line and that of S_v from the slop of the line, ϕ_{v0} and S_v are the measure of solute- solvent and solvent- solvent interaction respectively. ϕ_v apparent molar volume is positive and show sharp decrease with the increase of soap concentration below c_{mc} and slight increase above c_{mc} Fig.1 (A, B, C). ϕ_v , apparent molar volume is solute-solvent inter action, the positive value indicates strong solute- solvent inter action, a progress for overall structural order in alkanols. The soaps in alkanols behave actively as structure making agent. The decline in ϕ_v is as soaps molecules in dilute solution ionise considerably in to cation Rb⁺ and anion hydrocarbon which are encircled completely by a layer of solvent molecules, firmly bounded by Vander Walls' forces and dipol-dipol interaction which are oriented towards the ions. Change in overall structure is evident.

The apparent molar volume is extrapolated to zero soap concentration in the graph of ϕ_v vs \sqrt{C} , to get limiting apparent molar volume ϕ_v^0 i.e. as follows:

$$\phi_v = \phi_v^\infty = \lim (m \rightarrow 0) \phi_v$$

Table 1B: Parameters from ϕV vs \sqrt{C} graphs

Parameters	Rubidium soaps								
	Caprylate			Caprate			Laurate		
	Methanol	Butanol	Pentanol	Methanol	Butanol	Pentanol	Methanol	Butanol	Pentanol
$\phi_{v01} \text{cm}^3 \text{mol}^{-1}$	340.1	203.2	251.0	154	111	160	168.3	36.1	48.3
$\phi_{v02} \text{cm}^3 \text{mol}^{-1}$	71.0	12	199.7	59.8	104.5	159.8	59.8	1.1	7.9
$S_v \text{cm}^3 \text{mol}^{-3/2}$	-1.1	-0.84	-0.67	-0.34	-0.39	-0.31	-0.42	-0.071	-0.069

Massion equation is applicable to the system below cmc as well as above c mc, two values of ϕ_v^0 are possible ϕ_{v1}^0 below cmc and ϕ_{v2}^0 above cmc. It is a measure of solute- solvent interaction below cmc and its positive value with higher magnitude Table 1B verifies quantitatively strong solute –solvent interaction leading to change in overall structural order, The structure breaking effect of soap due to ion-solvent inter action between the molecules may be visualized. It is concluded that interaction of soap and solvent is higher below cmc whereas soap- soap interaction exists due to its low value. Experimental slop S_v has values, S_{v1} below c mc and S_{v2} above c mc. S_{v1} comes out negative pointing towards weak ion-ion inter action as a result of inter ionic penetration Table1B whereas, S_{v2} is positive. The S_{v2} may have positive value due to appreciable ionization of soap at higher concentration. The solute –solute inter action is appreciably strong as the concentration of soap in solution increases due to micellisztion process.

CONCLUSION

The rubidium soaps considerably ionize in solution to form anion hydrocarbon (caprylate, caprate, laurate) giving rise to hydrogen bonding favoring condition and resonance. Massion equation $\phi_v = \phi_{v0} + S_v \sqrt{C}$ is applicable to the dilute solution of soaps under investigation. The value of ϕ_{v1}^0 and ϕ_{v2}^0 apparent molar volume below cmc and above cmc depend on the chain length of the soaps. The positive value & deceasing character of apparent molar volume indicates strong ion-solvent interaction and shows structure breaking capacity of soaps in alkanols.

ACKNOWLEDGEMENTS

I am thankful to Prof. R.C. Kapoor, D.Sc., D. Phil., F.N. A.Sc., Head of Chemistry Department, University of Jodhpur, Jodhpur for his keen interest and valuable criticism throughout. I am thankful to Prof. Chanderdeep Tondon, Director Amity Institute of Biotechnology, 125, Noida. I am also thankful to prof. K. C. Upadhayay, FNA, FNASc, Advisor Amity Institute of Biotechnology, Noida.

CONFLICT OF INTEREST: None

REFERENCES

- [1] Burrows HD, Chimamkpam TO, Encarnacao T et al. (2010). Trivalent metal ion binding to surfactants and polyelectrolytes; A review. *J Surface Sci. Technol.* 26(3-4): 1-16
- [2]. Liu YB, Liu WQ, Meng HH (2007). Metal dicarboxylates as thermal stabilizers for PVC. *Degrad Stab.* 92 (8): 1565-71
- [3] Barbara M, Lacz A (2007). Thermal decomposition of cadmium succinate dehydrate. *J Therm. Anal and Calorimetry.* 88 (1): 295-99

- [4] Shukla M, Kumari S, Shukla RK (2010). Effect of chain length on acoustic behavior of gadolinium alkanoates in mixed organic solvents, *Acta Acustion*. 96: 63-67
- [5] Mehrotra KN, Anis (1997). Apparent molar volume and viscosity of zirconyl soaps, *J. colloid and surface*. 122 (1-3): 27-32
- [6] Pereira PRF, Valente JMA, Burrows DH (2010). Thermodynamic analysis of the interaction between trivalent metal ions and sodium dodecyl sulfate. *J Mol Liquids*. 156(1): 109-114
- [7] Sagar V, Sharma M, Sharma V. (2006). Viscosity of thorium soaps. *Asian J Chem*. 18(4): 3065-72
- [8]. Shukla M, Kumari S, Shukla RK (2011). Effect of temperature on micellization and thermodynamic behavior of gadolinium alkanoates in non-aqueous media, *J. Dispers Sci Tech*. 32, 1668-72
- [9] Sharma R, Tank P, Saxena M et al. (2008). Synthesis and characterisation of antifungal agents containing copper(II) soaps and derived from mustard and soyabean oil. *Tenside Surfactants Detergents*. 45(2): 87-92
- [10] Mehrotra KN, Shukla RK, Chauhan M (1996). Conductometric investigations of samarium valerate, caproate and caprylate in methanol. *J. Colloid and Surface*. 119: 67-72
- [11] Sharma V, Sharma M, Sagar V (2006). Viscometric investigations of uranyl soap solutions at different temperature. *J Indian Chem Soc*. 83: 1270 -72
- [12] Patil RS, Animesh RK (2003). Physicochemical properties of anionic-nonionic surfactant mixture: α -sulfonatomyristic acid methyl ester (MES)-nonaoxyethylene monododecyl ether (C12E9). *J of the Indian Chem Soc*. 80(4): 345-50
- [13] Mehrotra KN, Saroha SPS (1979). Studies on infrared spectra and thermal analysis of rubidium soaps. *J Indian Chem Soc*. LVI: 466-469



ISSN 2456-2580